

$U(1)$ invariant Membranes and Singularities

Jens Hoppe

Abstract. A formulation of $U(1)$ - symmetric classical membrane motions (preserving one rotational symmetry) is given, and reductions to systems of ODE's, as well as some ideas concerning singularities and integrability.

Let me start by giving some particular example(s) of a 2-dimensional (time-like, time periodic) extremal manifold $\mathbb{M}_2 \subset \mathbb{R}^{1,2}$ and 3-folds (singularities included) of vanishing mean curvature, in 4-dimensional Minkowski space ([1],[2]):

$$\mathbb{M}_2 := \left\{ \begin{pmatrix} t \\ x \\ y \end{pmatrix} \in \mathbb{R}^{1,2} \mid x \cos 4t - y \sin 4t = 8(x^2 + y^2)^2 - 3(x^2 + y^2) - \frac{3}{32}; x^2 + y^2 > \frac{1}{16} \right\},$$

$$\mathbb{M}_3^{(t)} := \{(x^\mu) \in \mathbb{R}^{1,3} \mid \mathcal{P}(x)\mathcal{P}(y)\mathcal{P}(t) = \mathcal{P}(z)\}, \quad (1)$$

$$\mathbb{M}_3^{(s)} := \{(x^\mu) \in \mathbb{R}^{1,3} \mid \mathcal{P}(x)\mathcal{P}(y)\mathcal{P}(z) = \mathcal{P}(t)\},$$

where $\mathcal{P} : \mathbb{R} \rightarrow \mathbb{R}_{\geq 1}$ is a (2ω) real periodic) elliptic Weierstrass function satisfying

$$\mathcal{P}'^2 = 4\mathcal{P}(\mathcal{P}^2 - 1), \quad \mathcal{P}(\omega) = 1.$$

$\mathbb{M}_3^{(s)}$ can be thought of as a (infinite assembly of) melting ice-block(s), as at $t = 0$ it covers the surface(s) of infinitely many cubes (each of size 2ω), packed together, for $0 < t < T = 2\omega$ instantaneously separating into infinitely many melting (round/ed) ice piece(-surface)s, each of which shrinks to a point ($t = T/2 = \omega$) and then growing again to become, at $T = 2\omega$, the surface of infinitely many ice-cubes, stacked together. $\overline{\mathbb{M}}_2$ (with the $x^2 + y^2 = \frac{1}{16}$ line of singularities included) gives the motion of a closed string of a fixed (heart-like) shape (with exactly one cusp - singularity, that arises when the radius r takes its minimum, $r = \frac{1}{4}$), rotating with constant angular velocity ($=4$) around the origin [6]. Before discussing time-like extremal 3-manifolds with one rotational symmetry embedded in $\mathbb{R}^{1,4}$, consider for a moment the general (parametric) description of $(M + 1)$ -dimensional extended

Based on talks given at the Banff Centre (July 2007) and the Institut Henri Poincare (March 2008).

manifolds $\mathbb{M}$ embedded in D -dimensional Minkowski-space: varying the volume-functional

$$\text{Vol}(\mathbb{M}) = \int d^{M+1}\varphi \sqrt{G} = S[x^\mu] \quad (2)$$

with respect to the embedding functions $x^\mu(\varphi^0, \dots, \varphi^M)$, $\mu = 0, \dots, D-1$, one obtains the parametric extremal volume equations,

$$\frac{1}{\sqrt{G}} \partial_\alpha (\sqrt{G} G^{\alpha\beta} \partial_\beta x^\mu) = 0, \quad (\mu = 0, \dots, D-1), \quad (3)$$

where $G_{\alpha\beta} := \partial_\alpha x^\mu \partial_\beta x^\nu \eta_{\mu\nu}$, $\eta_{\mu\nu} = \text{diag}(1, -1, \dots, -1)$, $G^{\alpha\beta} G_{\beta\gamma} = \delta_\gamma^\alpha$, $G = (-1)^M \det(G_{\alpha\beta})$, $\alpha, \beta = 0, \dots, M$.

In the gauge $\varphi^0 = x^0 (= t)$, i.e. $(x^\mu)^T = (t, \vec{x}(t, \varphi^1, \dots, \varphi^M))$, and also demanding

$$\sum_{i=1}^{D-1} \dot{x}_i \frac{\partial x_i}{\partial \varphi^r} = 0, \quad (r = 1, \dots, M), \quad (4)$$

eq. (3) $_{\mu=0}$ becomes a conservation law

$$\dot{\rho} = 0, \quad \rho := \sqrt{\frac{g}{1 - \dot{\vec{x}}^2}} \quad (5)$$

while the remaining ones, (3) $_{\mu=i}$, become the evolution equations for a M -dimensional time-dependent surface $\Sigma(t)$ moving in $\mathbb{R}^{D-1}$,

$$\ddot{x}_i = \frac{1}{\rho} \partial_r \left(\frac{g}{\rho} g^{rs} \partial_s x_i \right) \quad (i = 1, \dots, D-1); \quad (6)$$

g^{rs} respectively g are the inverse, respectively the determinant, of the M -dimensional metric on $\Sigma(t)$, g_{rs} , induced from $\mathbb{R}^{D-1}$. In [3], I also considered the equations

$$\ddot{\vec{x}} = (1 - \dot{\vec{x}}^2) g^{rs} \partial_{rs}^2 \vec{x}, \quad (7)$$

respectively

$$\ddot{\vec{x}} = \frac{1}{\rho^2} g g^{rs} \partial_{rs}^2 \vec{x}. \quad (8)$$

The following calculations show that (8) and (4) consistently imply (5) and (6):

$$\begin{aligned} \frac{1}{2} (1 - \dot{\vec{x}}^2) \frac{d}{dt} (\rho^2) &= \frac{1}{2} \dot{g} + g \frac{\dot{\vec{x}} \ddot{\vec{x}}}{1 - \dot{\vec{x}}^2} \\ &= \frac{1}{2} g g^{rs} \dot{g}_{rs} + \frac{g}{\rho^2 (1 - \dot{\vec{x}}^2)} (g g^{rs} \partial_{rs}^2 \vec{x}) \dot{\vec{x}} \\ &= g g^{rs} \partial_r \vec{x} \partial_s \dot{\vec{x}} + g g^{rs} \dot{\vec{x}} \partial_{rs}^2 \vec{x} = 0. \end{aligned} \quad (9)$$

In order to deduce that (8) implies (6) one may equivalently show that (in a parametrization which (8) and (4) hold) necessarily

$$\partial_r \left(\frac{g g^{rs}}{\rho} \right) \partial_s \vec{x} = 0 ; \quad (10)$$

multiplying by $\frac{\rho}{g} \partial_a \vec{x}$ one obtains

$$\begin{aligned} \frac{\rho}{g} \partial_a \left(\frac{g}{\rho} \right) + \partial_r (g^{rs}) g_{sa} &= \frac{1}{2} \frac{\partial_a g}{g} - \frac{\dot{\vec{x}} \partial_a \dot{\vec{x}}}{1 - \dot{\vec{x}}^2} - g^{rs} (\partial_{rs}^2 \vec{x} \partial_a \vec{x} + \partial_s \vec{x} \partial_{ra}^2 \vec{x}) \\ &= \frac{\ddot{\vec{x}} \partial_a \vec{x}}{1 - \dot{\vec{x}}^2} - g^{rs} \partial_{rs}^2 \partial_a \vec{x} = 0. \end{aligned} \quad (11)$$

Let us now restrict to the case of membranes ($M = 2$), as well as to $\Sigma(t)$ being rotationally symmetric, of the form

$$\vec{x}(t, \varphi^1 \varphi^2) = \begin{pmatrix} u_1(t, \varphi^1) \begin{pmatrix} \cos \varphi^2 \\ \sin \varphi^2 \end{pmatrix} \\ \vdots \\ u_N(t, \varphi^1) \begin{pmatrix} \cos \varphi^2 \\ \sin \varphi^2 \end{pmatrix} \end{pmatrix}. \quad (12)$$

Noting that (g_{rs}) , hence also g, g^{rs} , and ρ , will be independent of φ^2 , $\vec{e}(t, \varphi) := a \vec{u}(\frac{t}{a}, \varphi^1(\varphi))$, with $\varphi(\varphi^1)$, respectively $\varphi^1(\varphi)$ defined via $a^2 \frac{d\varphi}{d\varphi^1} = \rho(\varphi^1)$ (where $a^2 := \langle \rho \rangle$, the average value of ρ) equation(s) (6) may be written in the form

$$\ddot{\vec{e}} = \vec{e}^2 \vec{e}'' + 2(\vec{e} \dot{\vec{e}}) \dot{\vec{e}} - \dot{\vec{e}}^2 \vec{e}, \quad (13)$$

to be supplemented by the two constraints (cp. (4)/(5))

$$\dot{\vec{e}} \dot{\vec{e}} = 0, \quad (14)$$

$$\dot{\vec{e}}^2 + \vec{e}^2 \dot{\vec{e}}^2 = 1. \quad (15)$$

Note that, when $D = 5$ ($N = 2$), and as long as $\dot{\vec{e}}$ and $\vec{e}$ are linear independent, these 2 conditions (14) and (15) actually imply (13), - which can be seen by comparing the 4 equations obtained from differentiating (14) and (15) with respect to t and φ with the scalar product of (13) with $\dot{\vec{e}}$, respectively $\vec{e}$. Also note that (14) and (15), which may be thought of as evolution equations of a string moving in the plane, $\dot{\vec{e}} = v \vec{n}$ (with $\vec{n}$ being the unit normal to $\vec{e}$ and the magnitude of the normal velocity being $\sqrt{1 - \vec{e}^2 \dot{\vec{e}}^2}$), consistently reduce to a system of ODE's when making the Ansatz

$$\vec{e}(t, \varphi) = R(\omega t) \vec{f}(\varphi + ct), \quad (16)$$

where $R(\omega t)$ is given by

$$R(\omega t) = e^{\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \omega t} = \begin{pmatrix} \cos \omega t & \sin \omega t \\ -\sin \omega t & \cos \omega t \end{pmatrix} \in SO(2)$$

and the possible shapes $\vec{f} = r \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix}$ being determined by solving

$$\omega^2(u - c^2)^2 u'^2 = 4(\omega^2 u - 1)(c^2 - \omega^2 u^2)$$

in terms of elliptic integrals involving $u = r^2(\tilde{\varphi} := \varphi + ct)$, and $\frac{d\theta}{d\tilde{\varphi}} = \frac{\omega^2 u - 1}{c^2 - u} \frac{c}{\omega u}$

A more general approach to solving (14) and (15) results when writing $\vec{e}$ in polar form,

$$\vec{e}(t, \varphi) = r(t, \varphi) \begin{pmatrix} \cos \psi(t, \varphi) \\ \sin \psi(t, \varphi) \end{pmatrix}, \quad (17)$$

and the corresponding equations,

$$\begin{aligned} \dot{r}\dot{r} + r^2\dot{\psi}\dot{\psi} &= 0, \\ \dot{r}^2 + r^2\dot{\psi}^2 + r^2(\dot{r}^2 + r^2\dot{\psi}^2) &= 1 \end{aligned} \quad (18)$$

as

$$(r\dot{\psi} \pm r^2\dot{\psi})^2 + (\dot{r} \pm r\dot{r})^2 = 1. \quad (19)$$

Eq. (13), on the other hand, is consistent with the Ansatz (cp. [4] for the $N = 1$ case)

$$\vec{e} = t^a \vec{f}(t^b \varphi), \quad (a + b + 1 = 0), \quad (20)$$

yielding the system of ODE's

$$\vec{f}^2 \vec{f}'' + 2(\vec{f} \dot{\vec{f}}) \dot{\vec{f}} - \dot{\vec{f}}^2 = a(a-1)\vec{f} + (2-a)(a+1)s\vec{f} + (a+1)^2 s^2 \vec{f}'' \quad (21)$$

for $\vec{f}(s)$. For $a = -1$, eq. (21) follows from

$$H = \frac{1}{2} \frac{\vec{p}^2}{\vec{f}^2} - \vec{f}^2 \quad \text{resp.} \quad L = \frac{1}{2} \vec{f}^2 \dot{\vec{f}}^2 + \vec{f}^2. \quad (22)$$

While the system separates for general N , a particularly simple solution can be obtained for $N = 2$, for which

$$L = \frac{1}{2} r^2 (\dot{r}^2 + r^2 \dot{\theta}^2 + 2) \quad (23)$$

when using $\vec{f}^T = r(s) (\cos \theta(s), \sin \theta(s))$.

$$\frac{d}{ds} (r^4 \dot{\theta}) = 0, \quad \dot{\theta} = \frac{L}{r^4} \quad (24)$$

allows to eliminate $\dot{\theta}$,

$$L = \frac{1}{2} r^2 (\dot{r}^2 + \frac{1}{2} \frac{L^2}{r^4}) + r^2, \quad (25)$$

and

$$\frac{1}{2} r^2 \dot{r}^2 + \frac{1}{2} \frac{L^2}{r^4} - r^2 = E \quad (26)$$

to be constant, leading to the elliptic integral, also obtained by M. Trzetrzelewski (private communication), for $v = r^2$

$$\int ds = \pm \int \frac{v dv}{\sqrt{2v^3 + 2Ev^2 - L^2}}. \quad (27)$$

A light cone version of (13) with $t(=x^0) \rightarrow \tau := \frac{x^0+x^5}{2}$, i. e.

$$\vec{e} = \frac{2}{x^0 + x^5} \sqrt{v} \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix},$$

implying $4v = (x^0 + x^5)^2(x_1^2 + x_2^2 + x_3^2 + x_4^2)$ and obtaining $(x^0 - x^5) =: x_-$ via integrating $\partial_\varphi x_- = \partial_\tau \vec{x} \partial_\varphi \vec{x}$, leads to the extremal 3-folds

$$\mathbb{M}_{c,L} = \{ (x^\mu) \in \mathbb{R}^{1,5} \mid (x_0 - x_5)^2(x_1^2 + x_2^2 + x_3^2 + x_4^2 + x_0^2 - x_5^2) = c, \ x_1 x_4 - x_2 x_3 = 0, (*) \},$$

where the third condition (*) relating the 6 coordinates is obtained by eliminating v and θ , via (24) and (27) (with $8E = -3c$), from $(e_1)^2 = (x_1)^2 + (x_2)^2$ and $(e_2)^2 = (x_3)^2 + (x_4)^2$. Various other aspects, including the question of integrability of (14) and (15) are currently under investigation [5].

References

- [1] J. Hoppe, ‘Conservation Laws and Formation of Singularities in Relativistic Theories of Extended Objects’, in: ‘Non-linear Waves’, Vol.8 of the GAKUTO International Series, Tokyo 1996 (hep-th/9503069).
- [2] J. Hoppe, ‘Membranes and Matrix Models’, IHES/P/02/47, hep-th/0206192.
- [3] J. Hoppe, ‘Reformulating M-brane Equations’, Physics Letters B (in print).
- [4] J. Hoppe, ‘Some Classical Solutions of Relativistic Membrane Equations’, Physics Letters B 329 (1994) 10.
- [5] J. Hoppe, M. Trzetrzelewski and A.A. Zheltukhin; in preparation.
- [6] $\mathbb{M}_2$ arises as a special case ($\omega = 4, c = 3$) of rotating with angular velocity ω shapes given by $r \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix}$,

$$r^2(\tilde{\varphi}) = \frac{1}{2\omega^2} ((c^2 + 1) + (c^2 - 1) \cos k\tilde{\varphi}),$$

$$\frac{d\theta}{d\tilde{\varphi}} = \frac{1 - \omega^2 r^2}{1 - c^2} \frac{c\omega}{\omega^2 r^2} = \frac{kc}{4}(c^2 - 1) \frac{1 + \cos k\tilde{\varphi}}{1 + \frac{c^2-1}{2}(1 + \cos k\tilde{\varphi})},$$

$k = \frac{2\omega}{c^2-1}$; for $c = (2n+1) = c_n$ an odd integer, $\omega = 2n(n+1) = \omega_n$, ($k = 1$), the non-parametric description is $\omega_n x = \sqrt{w} \cos \theta = h_{n+1}(w := \omega^2 r^2)$, with h_{n+1} a polynomial (resp. $-\sqrt{w} \cos \left(c \arctan \sqrt{\frac{w-1}{c^2-w}} - \arctan \left(c \sqrt{\frac{w-1}{c^2-w}} \right) \right)$) satisfying

$$c^2(h^2 - w)(w - 1) = (h - 2wh')^2(w - c^2).$$

Acknowledgements

I would like to thank the organizers of the workshops ‘Quadrature Domains and Laplacian Growth in Modern Physics’, resp. ‘Microscopic Description of Singularities’, for their kind invitation, the Marie Curie Training Network ENIGMA and the Swedish Research Council for support, and P. Allen, L. Andersson, M. Bordemann, J. Eggers, B. Gustafsson, V. Moncrief, A. Restuccia, S. Theisen, M. Trzetrzelewski, and A. Zheltukhin for discussions.

Jens Hoppe
Department of Mathematics, Royal Institute of Technology
10044, Stockholm, Sweden